

PRACTITIONER PRE-PRINTED ORDERS
Ferric Derisomaltose (Monoferric®)
Intravenous (Iron) Therapy in Pregnancy for
Out Patient Use ONLY
(TRIAL Use Only)

To complete the order form, fill in required blanks and/or initial the appropriate boxes.
 Bulleted items will be initiated automatically.
 To delete orders, draw one line through the item and initial.

Allergies: See Allergy / Intolerance Record	Patient Weight _____ kg <input type="checkbox"/> Estimated <input type="checkbox"/> Actual
--	--

Posted Initial	ORDERS AND SIGNATURE	Page 1 of 2
----------------	-----------------------------	--------------------

	<p><u>Diagnosis and Inclusion Criteria</u></p> <ul style="list-style-type: none"> • Iron Deficiency Anemia (IDA) as per Clinical Trial Protocol • Criteria met as per Inclusion/Exclusion Criteria form, and enrolled in Iron isomaltoside/ferric derisomaltose versus Iron Sucrose Clinical Trial <p><u>Exclusion Criteria</u></p> <ul style="list-style-type: none"> • Does not meet inclusion criteria as per IDA Clinical Trial Protocol • Multiple gestation pregnancy • Significant comorbidities: (e.g. lung disease, clotting disorders, heart conditions, immunological conditions, cancer history or presently receiving cancer treatment) • IV iron treatment within the last 4 weeks
--	--

	<p><u>Investigations or Tests</u></p> <ul style="list-style-type: none"> • Blood work if not done in the last 2 weeks: <ul style="list-style-type: none"> • Iron studies (serum ferritin, TIBC, TSAT) • CBC, Reticulocyte count • B12 • Other: _____
--	---

	<p><u>Treatments</u></p> <ul style="list-style-type: none"> • Refer to Intravenous Iron Therapy Care Plan Insert (RQHR 1598)
--	--

	<p><u>Observation</u></p> <ul style="list-style-type: none"> • Baseline BP, HR prior to initiation of all doses, every 15 minutes during infusion(s) and at 30 minutes post infusion • Fetal Monitoring: Assess uterine activity and fetal heart rate using fetal Doptone®/doppler prior to initiating infusion and within 30 minutes post infusion NOTE: This monitoring is a minimum. If increase in fetal monitoring is required, or changes in uterine activity, contact obstetrical care provider STAT for guidance. • Observe peripheral IV site for pain, redness, or swelling prior to initiating infusion and q 15 - 30 minutes until infusion complete • Observe for signs of hypersensitivity reaction every 15 minutes during infusion and 30 minutes post infusion. (Refer to Appendix A: Iron Infusion Hypersensitivity Reactions Management Algorithm) • Observe for signs of anaphylactic reactions (i.e. diaphoresis, hypotension, collapse) every 15 minutes from initiation of infusion and for 30 minutes after the end of infusion • Out patients may be moved to a suitable observation area after completion of infusion
--	--

Date & Time	Practitioner Signature: Practitioner Name (printed):
------------------------	---


PRACTITIONER PRE-PRINTED ORDERS
Ferric Derisomaltose (Monoferric®)
Intravenous (Iron) Therapy in Pregnancy for
Out Patient Use ONLY
(TRIAL Use Only)

To complete the order form, fill in required blanks and/or initial the appropriate boxes.
 Bulleted items will be initiated automatically.
 To delete orders, draw one line through the item and initial.

Allergies: See Allergy / Intolerance Record	Patient Weight _____ kg <input type="checkbox"/> Estimated <input type="checkbox"/> Actual
--	---

Posted Initial	ORDERS AND SIGNATURE	Page 2 of 2
----------------	-----------------------------	--------------------

	<p>Medication</p> <ul style="list-style-type: none"> Stop all previous forms of oral iron Administer IV iron in a monitored resuscitative facility where physicians/code team available <p>NOTE: Pre-medications are not recommended. If patient has a history of sensitivity to oral iron medication or medications containing elemental iron, nursing staff shall notify obstetrical care provider for further pre-medication orders.</p> <p>The Pharmacy department will supply one of the following:</p> <ul style="list-style-type: none"> iron sucrose (Venofer®) 300 mg. Refer to CS-OS-1926 Intravenous Iron for Iron Deficiency Anemia (IDA) in ADULT Outpatients for further Practitioner orders. OR iron isomaltoside (alt name: ferric derisomaltose [Monoferric®]) 1000 mg
--	--

	<p>Anaphylaxis Management</p> <ul style="list-style-type: none"> Refer to Clinical Procedure CS-CP-0014 Anaphylaxis – Identification and Initial Treatment – Acute and Continuing Care Settings If severe anaphylaxis, obtain Serial Serum Tryptase (Roy Romanow Provincial Laboratory) as soon as possible following reaction <p style="text-align: right;">Tryptase Collection Procedure </p>
--	---

	<p>Discharge Planning</p> <ul style="list-style-type: none"> If stable after mild or moderate acute hypersensitivity reaction for 1 - 4 hours, may discharge patient home Provide requisition to patient upon discharge for post infusion bloodwork CBC, ferritin, TIBC, TSAT, to be completed at 30 days post-infusion
--	--

	<p>Study Coordinator/MBU Coordinator Use only:</p> <p>Coordinator Name: _____ Date Reviewed: MM/DD/YYYY: _____</p> <p>Randomization Group _____ Block _____</p> <p>Study Coordinator to identify and which of the following two medications will be received by the patient:</p> <p><input type="checkbox"/> iron sucrose (Venofer®) 300 mg in 250 mL NS. Refer to CS-OS-1926 Intravenous Iron for Iron Deficiency Anemia (IDA) in ADULT Outpatients</p> <p><input type="checkbox"/> iron isomaltoside (alt name: ferric derisomaltose [Monoferric®]) 1000 mg in 100 mL NS</p> <p>Faxed completed PPO to obstetrical care provider <input type="checkbox"/> Yes</p> <p>Confirm PPO received by infusing area <input type="checkbox"/> Yes</p> <p>Study ID assigned, patient notified of study ID <input type="checkbox"/> Yes</p> <p>Date(s) of appointment for iron infusion: _____ / _____ / _____</p> <p>Follow up Blood work due MM/DD/YYYY: _____</p>
--	---

Fax this PPO signed by obstetrician to Patient Blood Management Department at 306-766-3328 or email to patientbloodmanagement@saskhealthauthority.ca

Date & Time	Practitioner Signature:
	Practitioner Name (printed):